

BRIUMVITM

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)		t) PRE	PRESCRIBER INFORMATION			
Name:	DOB:	Pres	Prescriber Name:			
Address:			State License:			
City, State, Zip:		NIDI I	NPI #: DEA:			
Phone: Alt. Phone:		Addr	Address:			
Email: SS#:						
Gender: M F Weight:(lbs) Ht:					ax:	
Allergies:		Offic	e Contact:		Phone:	
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)						
Primary Insurance:			Secondary Insurance (If Applicable):			
Plan #:		Plan	Plan #:			
Group #:			Group #:			
RX Card (PBM):		RX C	RX Card (PBM):			
BIN: PCN:		BIN:		P	CN:	
CLINICAL INFORMATION						
☐ G35 MS (relapsing remitting) ☐ G	Other (Specify ICD-10 Code	e):				
ab Orders: Frequency:						
Has patient received/plans on receiving any live or live-attenuated vaccinations 4 weeks prior to starting Briumvi™ treatment? ☐ Yes ☐ No						
Has Patient received/plans on receiving any non-live vaccinations 2 weeks prior to starting Briumvi™ treatment? ☐ Yes ☐ No						
Has Quantitative Serum Immunoglobulin Screening been performed? \square Yes \square No (Serum Immunoglobulin levels:)						
Has patient received an HBV Screening? ☐ Yes ☐ No (Results: ☐ Negative ☐ Positive)						
BRIUMVI™ ORDERS						
Prescription type: ☐ New start ☐ Restar	t Continued therapy	Total Doses	Received:	Date of L	ast Injection/Infusion:	
Medication	Dose/Frequency			Refills		
	☐ First Infusion: 150 mg (1 vial)					
☐ Briumvi™ 150mg vial	\square Second Infusion: 450 mg (3 vials) (2 weeks after initial dose)			Refill:		
□ Briumvi 150mg viai	\square Subsequent Infusion: 450 mg (3 vials) once every 24 weeks					
	☐ Other:					
Pre-Medication	Route			Dose		
☐ Acetaminophen	□ РО	☐ 500mg	☐ 650mg	☐ 1000mg		
☐ Methylprednisolone (Solu-Medrol)	□IV	☐ 60mg	□ 100 mg	□mg		
☐ Diphenhydramine (Benadryl)	□ IV □ PO	☐ 25mg	☐ 50mg			
Other:						
INFUSION REACTION ORDERS						
Mild reaction protocol:						
☑ Diphenhydramine 25mg IV, one time, for pruritus.						
If symptoms worsen, see orders for moderate to severe reactions.						
Moderate reaction protocol:						
Acetaminophen 650mg PO, one time, for pyrexia or rigors						
☐ Diphenhydramine 50mg IV, one time, for pruritus or urticaria						
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms						

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



BRIUMVITM

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

		• •	`			
If symptoms worsen, see interventions for severe re	eactions					
Severe reaction protocol: (Call 911 if initiated):						
☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)						
oximes Diphenhydramine 50mg IV,one time, for respira	atory symptoms, edema, o	r anaphylaxis				
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis						
oxtimes Sodium Chloride 0.9% 500mL IV over 30-60 min	, one time, for cardiovascu	ılar symptoms				
oximes Epinephrine 0.3mg/0.3mL IM into mis-anterolat	teral aspect of thigh of ana	phylaxis, may repeat x1 in	5-15 minutes if symptoms are not resolved or			
worsen						
FLUSHING & LOCKING ORDERS						
Flushing Protocol (>66lbs/33kg)						
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:				
\boxtimes 0.9% Sodium Chloride 2-5mL IV flush before and	d after each infusion	\boxtimes 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw				
Locking Protocol (>66lbs/33kg)						
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-			
□ Heparin Sodium 10 units/mL 1mL IV final		nits/mL 3mL IV final	tunneled Catheter:			
flush post normal saline flush	flush post normal saline		☐ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush			
** May substitute Dextrose 5% in Water, or alternative,	for 0.9& Sodium Chloride, w	hen indicated due to incomp	atibility with medications bring infused			
SIGNATURE						
We hereby authorize Talis Healthcare LLC to provio medicine as prescribed in this referral.	de all supplies and additior	nal services (nursing/patier	nt training) required to provide and deliver the			

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

Prescriber Signature

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.