



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)		rt)	PRESCRIBER INFORMATION				
Name: DOB:			Prescriber Name: State License:				
Address:City, State, Zip:			NPI #: Tax ID:				
Phone: Alt. Phone:			Address:				
Email: SS#:			City, State, Zip:				
Gender: M F Weight:(lbs) Ht:			Phone: Fax:				
Allergies:			Office Contact: Phone:				
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)							
Primary Insurance:			Secondary Insurance (If Applicable):				
Plan #:			Plan #:				
Group #:							
			RX Card (PBM):				
BIN:	PCN:		BIN:	PCN:			
CLINICAL INFORMATION							
☐ G35 Multiple Sclerosis (MS) ☐ Other ICD-10 Code (Please Specify Diagnosis):							
DRUG ORDERS							
Prescription type: \square New start	\square Restart \square Continued therapy	Total [oses Received:	Date of Last	Injection/Ir	nfusion:	
Medication	Dose/Strength		Di	rections		Refills	
☐ Tysabri® (Natalizumab)	☐ 300mg/15ml Vial ☐ Other:		Infuse 300mg IV over 1 hour every 4 weeks Other:				
Pre-Medication	Dose/Strength		Directions				
☐ Acetaminophen	□ 500mg	□ Tak	Take 1-2 tablets PO prior to infusion or post-infusion as directed				
☐ Diphenhydramine	☐ 25mg IV/PO		Take 1 tablet PO prior to infusion or as directed OR				
☐ 50mg IV/PO ☐ Inject contents of 1 vial IV prior to infusion or as dir							
☐ Methylprednisolone	□ 40mg□ 100mg□ 125mg		☐ Inject contents of 1 vial IV prior to infusion or as directed☐ Other: Inject 100mg IV 30 minutes prior to infusion				
INFUSION REACTION ORDERS							
Mild reaction protocol:							
☑ Diphenhydramine 25mg IV, one time, for pruritus.							
If symptoms worsen, see orders for moderate to severe reactions.							
Moderate reaction protocol:							
Acetaminophen 650mg PO, one time, for pyrexia or rigors							
□ Diphenhydramine 50mg IV, or □							
	, one time, for respiratory or neuro	logic syn	nptoms				
If symptoms worsen, see interventions for severe reactions							

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.





Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Date: ___

Severe reaction protocol: (Call 911 if initiated):							
☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)							
☑ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis							
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis							
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms							
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or							
worsen							
FLUSHING & LOCKING ORDERS							
Flushing Protocol (>66lbs/33kg)							
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:					
\boxtimes 0.9% Sodium Chloride 2-5mL IV flush before and	d after each infusion	☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw					
Locking Protocol (>66lbs/33kg)			, 4.6.1.				
PIV and Midline: PICC:		Implanted Port, Tunneled Catheter, and Non-					
	⋈ Heparin Sodium 10 units/mL 3mL IV final		tunneled Catheter:				
flush post normal saline flush flush post normal saline		flush	$\ oxed{oxed}$ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush				
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused							
SIGNATURE							
We hereby authorize Talis Healthcare LLC to provio medicine as prescribed in this referral.	de all supplies and additior	nal services (nursing/patier	nt training) required to provide and deliver the				

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

Prescriber Signature

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.