



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMAT	ION (Co	omplete or F	ax Existing Char	t) PRESCRIBER INFORMATION			
PATIENT INFORMATION (Complete or Fax Existing Char							
	Name: DOB:						
	Address:City, State, Zip:				State License: Tax ID:		
							
Phone: Alt. Phone: Email: SS#:							
Gender:							
Allergies:				Office Contact:			
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)							
Primary Insurance:				Secondary Insurance (If Applicable):			
Plan #:							
Group #:							
RX Card (PBM):							
	PCN:						
CLINICAL INFORMAT							
		fy Diagnosis):					
,	•						
MG-ADL* score: Has the patient received Meningitis vaccination(s)? ☐ Yes ☐ No Date of vaccination(s): ☐ Please check this box if the patient has declined vaccination Reason:							
•			-	yes, Reason/Date:			
☐ Please check to confirm: The patient is enrolled in the SOLIRIS REMS program; The patient has been counseled about the risks of meningococcal							
infection; The patient has received information and a Patient Safety Card about the symptoms and risks of meningococcal infection.							
SOLIRIS® ORDERS							
Prescription type: New	escription type: New start Restart Continued therapy		Total Doses Received:	otal Doses Received: Date of Last Injection/Infusion:			
Medication	St	rength		Dose/Frequency		Refills	
				mg IV every			
☐ Soliris® (eculizumab)	☐ 300ı	mg/30mL	☐ Maintenance d	e dose: mg IV every weeks.			
			Other:				
Pre-Medication	Dose/Str		/Strength	Directions			
☐ Acetaminophen		□ 500mg		\square Take 1-2 tablets PO prior to infusion or post-infusion as directed			
☐ Diphenhydramine	☐ 25mg IV/PO ☐ 50mg IV/PO		0	☐ Take 1 tablet PO prior to infusion or as directed OR			
Dipliciniyaraninic			0	☐ Inject contents of 1 vial IV prior to infusion or as directed			
☐ Methylprednisolone	☐ 40mg ☐ 100mg ☐ 125mg		L00mg	☐ Inject contents of 1 vial IV prior to infusion or as directed			
			\Box Other: Inject 100mg IV 30 minutes prior to infusion				
INFUSION REACTION ORDERS							
Mild reaction protocol:							
☑ Diphenhydramine 25mg IV, one time, for pruritus.							
If symptoms worsen, see orders for moderate to severe reactions.							

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



SOLIRIS®

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Date: ___

Moderate reaction protocol:								
☐ Acetaminophen 650mg PO, one time, for pyrexia or rigors								
☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria								
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms								
If symptoms worsen, see interventions for severe reactions								
Severe reaction protocol: (Call 911 if initiated):								
☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)								
☑ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis								
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis								
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms								
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or								
worsen								
FLUSHING & LOCKING ORDERS								
Flushing Protocol (>66lbs/33kg)								
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:						
\boxtimes 0.9% Sodium Chloride 2-5mL IV flush before an	d after each infusion	☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mLIV flush after infusion/lab draw						
Locking Protocol (>66lbs/33kg)								
PIV and Midline:	dline: PICC:		Implanted Port, Tunneled Catheter, and Non-					
☐ Heparin Sodium 10 units/mL 1mL IV final	⊠ Heparin Sodium 10 ur	nits/mL 3mL IV final	tunneled Catheter:					
flush post normal saline flush	flush post normal saline	flush						
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused								
SIGNATURE								
We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.								

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

Prescriber Signature

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.