



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Com	plete or Fax Existing Cha	rt) PRESCRIBER INFORMATION	PRESCRIBER INFORMATION			
Name:	DOB:	Prescriber Name:				
Address:						
City, State, Zip:		I NIDLU. DEA.				
Phone: Al		l - · ·	Address:			
Email:	SS#:		City, State, Zip:			
Gender: ☐ M ☐ F Weight:	(lbs) Ht:					
Allergies:		Office Contact:	Office Contact: Phone:			
INSURANCE INFORMATION –	AND – Send a copy of the	patient's prescription/insurance card	(front & back)			
Primary Insurance:		Secondary Insurance (If Applicable):				
Plan #:		Plan #:				
Group #:						
RX Card (PBM):						
BIN:						
CLINICAL INFORMATION						
Primary ICD-10 (Please Specify Diagn	osis):	Secondary ICD-10 (Please Specify Diagno	Secondary ICD-10 (Please Specify Diagnosis):			
Tertiary ICD-10 (Please Specify Diagn	osis):					
Is the patient on iron, folate and/or v	vitamin B12 therapy? ☐ Yes ☐	\square No \square Is the patient on dialysis? \square Yes \square !	No			
Has patient received any ESA therapy	y? □ Yes □ No If yes, how	many weeks of ESA therapy has the patient com	pleted? weeks			
Patient's hemoglobin (Hgb) level:	g/dL					
ARANESP® ORDERS						
Prescription type: ☐ New start ☐	Restart Continued therapy	Total Doses Received: Date of	Last Injection/Infusion:			
Medication		Dose/Frequency	Refills			
☐ Aranesp® (darbepoetin alfa) Singl	e ☐ 25 mcg ☐ 40 m	ncg \square 60 mcg \square 100 mcg				
Dose Vials	☐ 150 mcg ☐ 200	mcg	Refills:			
☐ Aranesp® (darbepoetin alfa) Single Dose Prefilled Syringe	e □ Weekly □ Ever	y 2 Weeks Other:				
Special Instructions:						
Pre-Medication	Dose/Strength	Direction	ns			
☐ Acetaminophen	□ 500mg	☐ Take 1-2 tablets PO prior to infusion or po	Take 1-2 tablets PO prior to infusion or post-infusion as directed			
Dish salsadasasia s	☐ 25mg IV/PO	☐ Take 1 tablet PO prior to infusion or as directed OR				
☐ Diphenhydramine	☐ 50mg IV/PO	☐ Inject contents of 1 vial IV prior to infusion or as directed				
☐ Methylprednisolone	☐ 40mg ☐ 125mg	☐ Inject contents of 1 vial IV prior to infusion	Inject contents of 1 vial IV prior to infusion or as directed			
INFUSION REACTION ORDERS						
Mild reaction protocol:						
☐ Diphenhydramine 25mg IV, one time, for pruritus.						
If symptoms worsen, see orders for moderate to severe reactions.						
Moderate reaction protocol:						

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



ARANESP®

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Date: __

_				_	
XI.	Acetaminophen	650mg PO	one time	for nurevia	or rigors

- ☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria
- ☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms

If symptoms worsen, see interventions for severe reactions

Severe reaction protocol: (Call 911 if initiated):

- ☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)
- ☑ Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis

Prescriber Signature

 ☑ Methylprednisolone 125mg IV, one ti ☑ Sodium Chloride 0.9% 500mL IV over ☑ Epinephrine 0.3mg/0.3mL IM into mi worsen 	30-60 min, one time, for o	cardiovascular symp	• •		
FLUSHING & LOCKING ORDERS					
Flushing Protocol (>66lbs/33kg)					
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:			
\boxtimes 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion		\boxtimes 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw			
Locking Protocol (>66lbs/33kg)					
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-tunneled Catheter:		
☑ Heparin Sodium 10 units/mL 1mL IV final flush post normal saline flush	☑ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush		☑ Heparin Sodium 100 units/mL 3-5mL IV final flush post norm saline flush		
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused					
SIGNATURE					
We hereby authorize Talis Healthcare LL	C to provide all supplies a	nd additional servic	es (nursing/patient training) required to provide and deliver the		

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.