



PATIENT INFORMATION (Complete or Fax Existing Chart) PRESCRIBER INFORMATION

Name: _____ DOB: _____ Address: _____ City, State, Zip: _____ Phone: _____ Alt. Phone: _____ Email: _____ SS#: _____ Gender: <input type="checkbox"/> M <input type="checkbox"/> F Weight: _____ (lbs) Ht: _____ Allergies: _____	Prescriber Name: _____ State License: _____ NPI #: _____ Tax ID: _____ Address: _____ City, State, Zip: _____ Phone: _____ Fax: _____ Office Contact: _____ Phone: _____
---	--

INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)

Primary Insurance: _____ Plan #: _____ Group #: _____ RX Card (PBM): _____ BIN: _____ PCN: _____	Secondary Insurance (If Applicable): _____ Plan #: _____ Group #: _____ RX Card (PBM): _____ BIN: _____ PCN: _____
--	--

CLINICAL INFORMATION

Please Select Diagnosis:

G30.0 Alzheimer's disease with early onset
 G30.1 Alzheimer's disease with late onset
 G30.8 Other Alzheimer's disease
 G30.9 Alzheimer's disease, unspecified
 G31.84 Mild cognitive impairment, so stated
 Other: _____

Prescriber must indicate the following requirements have been met to confirm diagnosis and that Patient has evidence of AD neuropathology and has been assessed for baseline ARIA risk via MRI:

Amyloid pathology confirmed via:
 Amyloid PET Scan CSF analysis Blood plasma Date: _____ Result: Amyloid Positive Amyloid Negative

Recent MRI obtained prior to initiating Kisunla™ (including FLAIR and T2/GRE or SWI) to assess ARIA risk
 Prescriber has verified that this Patient does not have evidence of prior ARIA-H Date: _____

Completion of cognitive assessment type:
 MMSE MoCA CDR Other: _____ Date: _____

Completion of functional assessment type:
 FAQ FAST Other: _____ Date: _____

Completion of CMS approved CED registry (only required for Patients with Medicare) ClinicalTrials.gov Registry Number: NCT _____
 CED Submission Date: _____ Submission Number (if applicable): _____

****Note: MRIs must be obtained prior to initial infusion to assess ARIA risk. During treatment, conduct an ARIA monitoring MRI before Infusions 2, 3, 4 and 7 and if symptoms consistent with ARIA occur.**

LEQEMBI® ORDERS

Prescription type: New start Restart Continued therapy Total Doses Received: _____ Date of Last Injection/Infusion: _____

Medication	Dose/Frequency	Refills
<input type="checkbox"/> Leqembi® (lecanemab-irmb) 500 mg/5 mL (100 mg/mL) <input type="checkbox"/> Leqembi® (lecanemab-irmb) 200 mg/2 mL (100 mg/mL)	<input type="checkbox"/> 10 mg/kg intravenous infusion over approximately one hour, once every two weeks. <input type="checkbox"/> Other: _____	_____

Pre-Medication	Dose/Strength	Directions
<input type="checkbox"/> Acetaminophen	<input type="checkbox"/> 500mg	<input type="checkbox"/> Take 1-2 tablets PO prior to infusion or post-infusion as directed
<input type="checkbox"/> Diphenhydramine	<input type="checkbox"/> 25mg IV/PO <input type="checkbox"/> 50mg IV/PO	<input type="checkbox"/> Take 1 tablet PO prior to infusion or as directed OR Inject contents of 1 vial IV prior to infusion or as directed
<input type="checkbox"/> Methylprednisolone	<input type="checkbox"/> 40mg <input type="checkbox"/> 100mg <input type="checkbox"/> 125mg	<input type="checkbox"/> Inject contents of 1 vial IV prior to infusion or as directed <input type="checkbox"/> Other: Inject 100mg IV 30 minutes prior to infusion

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



<input type="checkbox"/>		
--------------------------	--	--

INFUSION REACTION ORDERS

Mild reaction protocol:
 Diphenhydramine 25mg IV, one time, for pruritus.
If symptoms worsen, see orders for moderate to severe reactions.

Moderate reaction protocol:
 Acetaminophen 650mg PO, one time, for pyrexia or rigors
 Diphenhydramine 50mg IV, one time, for pruritus or urticaria
 Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms
If symptoms worsen, see interventions for severe reactions

Severe reaction protocol: (Call 911 if initiated):
 Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)
 Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
 Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
 Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms
 Epinephrine 0.3mg/0.3mL IM into mid-antrolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen

FLUSHING & LOCKING ORDERS

Flushing Protocol (>66lbs/33kg)	
PIV and Midline: <input checked="" type="checkbox"/> 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion	Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter: <input checked="" type="checkbox"/> 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw

Locking Protocol (>66lbs/33kg)		
PIV and Midline: <input checked="" type="checkbox"/> Heparin Sodium 10 units/mL 1mL IV final flush post normal saline flush	PICC: <input checked="" type="checkbox"/> Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush	Implanted Port, Tunneled Catheter, and Non-tunneled Catheter: <input checked="" type="checkbox"/> Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush

**** May substitute Dextrose 5% in Water, or alternative, for 0.9% Sodium Chloride, when indicated due to incompatibility with medications being infused**

SIGNATURE

We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

X _____ Date: _____
 Prescriber Signature

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.