



## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

| PATIENT INFORMATION (Complete or Fax Existing Chart)   |  | t) PRESCRIBER I   | PRESCRIBER INFORMATION   |         |  |
|--|--|---|--|---------|--|
| Name:  |  | State License:  | Prescriber Name:   |         |  |
| INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)   |  |   |  |         |  |
| Primary Insurance:   |  | Plan #:<br>Group #:<br>RX Card (PBM): _   | Plan #: Group #: RX Card (PBM):  |         |  |
| CLINICAL INFORMATION   |  |   |  |         |  |
| □ E88.01 Alpha-1 antitrypsin deficiency □ Other ICD-10 code (Please Specify Diagnosis):  |  |   |  |         |  |
| GLASSIA® ORDERS  Prescription type:   New start   Restart   Continued therapy Total Doses Received:   Date of Last Injection/Infusion:   Date of Last Injection/Infusion:   New Start   Restart   Restart   Date of Last Injection/Infusion:   Date of Last Injection |  |   |  |         |  |
| Medication   | □ Restart □ Continued therapy                              | Dose/Frequency  |  | Refills |  |
| ☐ Glassia® [Alpha1-Proteinase Inhibitor (Human)]   | ☐ Infuse 60mg per kg (+/- 10%) nearest vial size) ☐ Other: | Infuse 60mg per kg (+/– 10%) intravenously weekly (where clinically appropriate, round to the earest vial size) |  |         |  |
| Pre-Medication   | Dose/Strength  |   | Directions   |         |  |
| ☐ Acetaminophen  | □ 500mg  | ☐ Take 1-2 tablets PO p   | Take 1-2 tablets PO prior to infusion or post-infusion as directed   |         |  |
| ☐ Diphenhydramine  | ☐ 25mg IV/PO<br>☐ 50mg IV/PO                               | •   | Take 1 tablet PO prior to infusion or as directed OR Inject contents of 1 vial IV prior to infusion or as directed |         |  |
| ☐ Methylprednisolone   | ☐ 40mg ☐ 125mg   | ☐ Inject contents of 1 vi   | Inject contents of 1 vial IV prior to infusion or as directed  |         |  |
|  |  |   |  |         |  |
| INFUSION REACTION ORDERS   |  |   |  |         |  |
| Mild reaction protocol:  ☑ Diphenhydramine 25mg IV, one time, for pruritus.  If symptoms worsen, see orders for moderate to severe reactions.  Moderate reaction protocol:  ☑ Acetaminophen 650mg PO, one time, for pyrexia or rigors  ☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria  ☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms   |  |   |  |         |  |

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.





## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

If symptoms worsen, see interventions for severe reactions Severe reaction protocol: (Call 911 if initiated): ☐ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%) ☑ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis ☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms ☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen **FLUSHING & LOCKING ORDERS** Flushing Protocol (>66lbs/33kg) PIV and Midline: Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter: □ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion ☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw Locking Protocol (>66lbs/33kg) PIV and Midline: PICC: Implanted Port, Tunneled Catheter, and Nontunneled Catheter: ☐ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush flush post normal saline flush ⋈ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush \*\* May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused **SIGNATURE** We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

**Prescriber Signature** 

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.