



## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)		PRESCRIBER INFORMATION	PRESCRIBER INFORMATION		
Name: DOB: Address: City, State, Zip: Alt. Phone:		State License: Tax ID:	Prescriber Name:		
Email:		City, State, Zip:			
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)					
Plan #: Group #: RX Card (PBM):	PCN:	Plan #: Group #: RX Card (PBM):			
CLINICAL INFORMATION					
☐ J45.50 Severe persistent asthma, uncomplicated ☐ J45.51 Severe persistent asthma with (acute) ☐ Other:					
FASENRA® ORDERS					
Prescription type: ☐ New start	☐ Restart ☐ Continued therapy	Total Doses Received: Date of Last	Injection/Infusion:		
Medication	Do	ose/Frequency	Frequency Refills		
☐ Fasenra® (benralizumab) 30mg/mL	<ul> <li>□ Starter Dose: Inject 30mg under doses, followed by once every 8</li> <li>□ Maintenance Dose: Inject 30mg</li> <li>□ Other:</li> </ul>		ks subsequently.		
Pre-Medication	Dose/Strength	Directions			
☐ Acetaminophen	□ 500mg	$\square$ Take 1-2 tablets PO prior to infusion or post-in	ake 1-2 tablets PO prior to infusion or post-infusion as directed		
☐ Diphenhydramine	☐ 25mg IV/PO ☐ 50mg IV/PO	'	ake 1 tablet PO prior to infusion or as directed OR nject contents of 1 vial IV prior to infusion or as directed		
☐ Methylprednisolone	☐ 40mg ☐ 100mg ☐ 125mg	'	nject contents of 1 vial IV prior to infusion or as directed ther: Inject 100mg IV 30 minutes prior to infusion		
INFUSION REACTION ORDERS					
Mild reaction protocol:  ☑ Diphenhydramine 25mg IV, one time, for pruritus.  If symptoms worsen, see orders for moderate to severe reactions.  Moderate reaction protocol:					

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.





Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Date: \_\_\_\_\_

☑ Acetaminophen 650mg PO, one time, for pyrexia or rigors						
☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria						
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms						
If symptoms worsen, see interventions for severe reactions						
Severe reaction protocol: (Call 911 if initiated):						
☐ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)						
☐ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis						
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis						
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms						
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or						
worsen						
FLUSHING & LOCKING ORDERS						
Flushing Protocol (>66lbs/33kg)						
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:				
$\boxtimes$ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion		☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw				
Locking Protocol (>66lbs/33kg)						
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-			
□ Heparin Sodium 10 units/mL 1mL IV final			tunneled Catheter:			
flush post normal saline flush	flush post normal saline flush		⊠ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush			
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused						
SIGNATURE						
We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.						

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

**Prescriber Signature** 

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.