



## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)			PRESCRIBER INFORMATION			
Name: DOB:  Address:  City, State, Zip: Alt. Phone:  Phone: Alt. Phone:  Email: SS#:  Gender:		Prescriber Name:				
INSURANCE INFORMATIO	N – AND – Send a copy of the	patient'	's prescription/insurance	cards (front 8	& back)	
Plan #: Group #: RX Card (PBM):	PCN:		Secondary Insurance (If Application #: Group #: RX Card (PBM): BIN:			
CLINICAL INFORMATION						
☐ G36.0 Neuromyelitis optica ☐ Other ICD-10/Diagnosis:						
UPLIZNA® ORDERS						
Prescription type:  New start Restart Continued therapy Total Doses Received: Date of Last Injection/Infusion:						
Medication		Dose/Fr	Dose/Frequency Qty/Refills			
Uplizna® (inebilizumab injection)				Quantity: Refills:		
Pre-Medication	Dose/Strength	<del></del>	Directions			
☐ Acetaminophen ☐ Diphenhydramine	☐ 500mg ☐ 25mg IV/PO ☐ 50mg IV/PO	□ Take	ke 1-2 tablets PO prior to infusion or post-infusion as directed ke 1 tablet PO prior to infusion or as directed OR ect contents of 1 vial IV prior to infusion or as directed			
☐ Methylprednisolone	☐ 40mg ☐ 100mg ☐ 125mg	1	ject contents of 1 vial IV prior to infusion or as directed ther: Inject 100mg IV 30 minutes prior to infusion			
		<u></u>				
Mild reaction protocol:  ☑ Diphenhydramine 25mg IV,  If symptoms worsen, see orders  Moderate reaction protocol:  ☑ Acetaminophen 650mg PO,	one time, for pruritus. s for moderate to severe reactions.					

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.





## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

oximes Diphenhydramine 50mg IV, one time, for prurit	us or urticaria						
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms							
If symptoms worsen, see interventions for severe reactions							
Severe reaction protocol: (Call 911 if initiated):							
☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)							
☑ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis							
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis							
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms							
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or							
worsen							
FLUSHING & LOCKING ORDERS							
Flushing Protocol (>66lbs/33kg)							
PIV and Midline:	Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:						
☑ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion		☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL					
	IV flush after infusion/lab draw						
Locking Protocol (>66lbs/33kg)							
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-				
☐ Heparin Sodium 10 units/mL 1mL IV final	☐ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush		tunneled Catheter:				
flush post normal saline flush			☑ Heparin Sodium 100 units/mL 3-5mL IV final				
	flush post normal saline flush						
** May substitute Dextrose 5% in Water, or alternative,	, for 0.9& Sodium Chloride, w	hen indicated due to incon	npatibility with medications bring ir	ıfused			
SIGNATURE							
We hereby authorize Talis Healthcare LLC to prov medicine as prescribed in this referral.	ide all supplies and additic	onal services (nursing/pat	ient training) required to provid	e and deliver the			
X			Date:	Prescriber			

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

Signature

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.