



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)		PRESCRIBER INFORMATION	PRESCRIBER INFORMATION	
Name:	DOB:	Prescriber Name:		
Address:		State License:		
City, State, Zip:		NPI #:Tax ID: _	NPI #: Tax ID:	
Phone: Alt. Phone:			Address:	
Email:	SS#:	City, State, Zip:		
Gender: ☐ M ☐ F Weight:	(lbs) Ht:	Phone: F	ax:	
Allergies:		Office Contact:	Phone:	
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)				
Primary Insurance:		Secondary Insurance (If Applicable):		
Plan #:				
Group #:				
RX Card (PBM):		RX Card (PBM):		
BIN:				
CLINICAL INFORMATION				
ICD-10 Code (Required):	ICD-10 Description	on:		
Baseline Liver Enzymes, Including Bilir	ubin (Results):			
Date of Negative TB Test:				
OMVOH™ ORDERS				
Prescription type: New start Restart Continued therapy Total Doses Received: Date of Last Injection/Infusion:				
Medication	Dose/Frequency		Qty/Refills	
☐ Induction Dosing: 300 mg / 15 mL single dose vial	300 mg by intravenous infusion at Weeks 0, 4, and 8		Quantity:	
☐ Maintenance Dosing: Omvoh™ 2 x 100 mg/mL given as two consecutive SQ injections	2 x 100 mg by subcutaneous injection every 4 weeks (thereafter)		Quantity:	
Pre-Medication	Dose/Strength	Directions		
☐ Acetaminophen	☐ 500mg	Take 1-2 tablets PO prior to infusion or post-infusion as directed		
·	☐ 25mg IV/PO	☐ Take 1 tablet PO prior to infusion or as directed OR		
☐ Diphenhydramine	☐ 50mg IV/PO	☐ Inject contents of 1 vial IV prior to infusion or as directed		
☐ Methylprednisolone	☐ 40mg ☐ 100mg	☐ Inject contents of 1 vial IV prior to infusion	Inject contents of 1 vial IV prior to infusion or as directed	
	☐ 125mg	Other: Inject 100mg IV 30 minutes prior to infusion		
INFUSION REACTION ORDERS				
Mild reaction protocol:				
☑ Diphenhydramine 25mg IV, one time, for pruritus.				
If symptoms worsen, see orders for moderate to severe reactions.				
Moderate reaction protocol:				
☐ Acetaminophen 650mg PO, one time, for pyrexia or rigors				
☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria				
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms				

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

If symptoms worsen, see interventions for severe reactions

Severe reaction protocol: (Call 911 if initiated):

- ☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)
- ☑ Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
- ☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
- ☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms
- ⊠ Enipenhrine 0.3mg/0.3ml. IM into mis-anterplateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or

worsen					
FLUSHING & LOCKING ORDERS					
Flushing Protocol (>66lbs/33kg)					
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:			
☑ 0.9% Sodium Chloride 2-5mL IV flush before an	d after each infusion	\boxtimes 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw			
Locking Protocol (>66lbs/33kg)					
PIV and Midline: ☑ Heparin Sodium 10 units/mL 1mL IV final	PICC: ⊠ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush		Implanted Port, Tunneled Catheter, and Non- tunneled Catheter:		
flush post normal saline flush					
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused					
SIGNATURE					
We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.					
x			Date:		
Prescriber Sig	gnature				

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.