



## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)			PRESCRIBER INFORMATION				
Name: DOB:			Prescriber Name:				
Address:			State License:				
City, State, Zip:			NPI #: Tax ID:				
Phone: Alt. Phone:							
Email: SS#:							
Gender:   M   F   Weight:(lbs)   Ht:							
Allergies:			Office Contact:		Phone:		
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)							
Primary Insurance:			Secondary Insurance (If Applicable):				
Plan #:			Plan #:				
Group #:			Group #:				
RX Card (PBM):		RX Card (PBM):					
BIN:	PCN:		BIN:	PCN:			
CLINICAL INFORMATION							
☐ K51.919 Ulcerative colitis, unspecified with unspecified complications ☐ K50.90 Crohn's disease, unspecified without complications ☐ Other ICD-10 code (Please Specify Diagnosis):							
Has patient received a TB test?					] Midline		
ENTYVIO® ORDERS							
Prescription type:  New start Restart Continued therapy Total Doses Received: Date of Last Injection/Infusion:							
Medication	Dose/Frequency Refi			Refills			
☐ Entyvio® (vedolizumab)	<ul> <li>□ Infuse 300mg intravenously at weeks 0, 2, 6 and then every 8 weeks thereafter.</li> <li>□ Infuse 300mg intravenously every 8 weeks.</li> <li>□ Other:</li></ul>						
Pre-Medication	Dose/Strength	Directions					
☐ Acetaminophen	□ 500mg	☐ Tak	☐ Take 1-2 tablets PO prior to infusion or post-infusion as directed		ion as directed		
☐ Diphenhydramine	☐ 25mg IV/PO ☐ 50mg IV/PO		☐ Take 1 tablet PO prior to infusion or as directed OR ☐ Inject contents of 1 vial IV prior to infusion or as directed				
☐ Methylprednisolone	☐ 40mg ☐ 125mg	☐ Inject contents of 1 vial IV prior to infusion or as directed					
INFUSION REACTION ORDERS							
If symptoms worsen, see intervent Severe reaction protocol: (Call 91	e time, for pyrexia or rigors e time, for pruritus or urticaria one time, for respiratory or neuro tions for severe reactions 1 if initiated):				(>0.50%)		
☐ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)							

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



☑ Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis



## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

☑ Methylprednisolone 125mg IV, one time, for res	spiratory symptoms, eden	na, or anaphylaxis				
⊠ Sodium Chloride 0.9% 500mL IV over 30-60 min	, one time, for cardiovasc	ular symptoms				
oximes Epinephrine 0.3mg/0.3mL IM into mis-anterolat	eral aspect of thigh of and	aphylaxis, may repeat x1 ir	n 5-15 minutes if symptoms are not resolved or			
worsen						
Flushing Protocol (>66lbs/33kg)						
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:				
□ 0.9% Sodium Chloride 2-5mL IV flush before and	d after each infusion	☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL				
		IV flush after infusion/lab draw				
Locking Protocol (>66lbs/33kg)						
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-			
☐ Heparin Sodium 10 units/mL 1mL IV final	☑ Heparin Sodium 10 units/mL 3mL IV final		tunneled Catheter:			
flush post normal saline flush	flush post normal saline	flush				
			flush post normal saline flush			
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused						
SIGNATURE						
We hereby authorize Talis Healthcare LLC to provio medicine as prescribed in this referral.	de all supplies and addition	nal services (nursing/patie	nt training) required to provide and deliver the			
X			Date:			
Prescriber Sig	nature					

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.